arXiv:1210.4326v1 [math.CA] 16 Oct 2012

ON THE OPTIMALITY OF A THEOREM OF KOKSMA

MICHEL J.G. WEBER

Abstract. We show the optimality of a result proved by Koksma in 1953, on the convergence almost everywhere of averages of dilated functions on the circle.

1. Introduction-Result

Throughout this Note, let $\mathbb{T}=\mathbb{R}/\mathbb{Z}\simeq[0,1[$ denotes the circle endowed with the normalized Lebesgue measure λ and $L^p=L^p(\lambda)$. Let also $e(x)=\exp(2i\pi x)$, $e_n(x)=e(nx)$, $n\in\mathbb{Z}$.

In 1953, Koksma published in [4] the following result.

Theorem 1.1. *Let* $g\in L^2$, $g\sim\sum_{k\in\mathbb{Z}}a_ke_k$ *be such that*

$$\sum_{k=1}|a_k|^2\sigma_{-1}(k)<\infty, \tag{1.1}$$

where $\sigma_{-1}(k)=\sum_{d|k}\frac{1}{d}$. *Then for almost all* x,

$$\frac{1}{n}\sum_{k=1}^{n}g(kx)\ \to\ \int_0^1 g(t)\mathrm{d}t,$$

as n *tends to infinity.*

This theorem is quite sharp. It improves a previous result in [3], where Koksma established the sufficiency of the coefficient condition

$$\sum_{k=1}|a_k|^2(\log\log k)^3<\infty. \tag{1.2}$$

The function $\sigma_{-1}(x)$ is multiplicative ($\sigma_{-1}(mn)=\sigma_{-1}(m)\sigma_{-1}(n)$ if m,n are coprime), and by Gronwall's estimate ([2] p.119–122),

$$\limsup_{x\to\infty}\frac{\sigma_{-1}(x)}{\log\log x}=e^{\lambda}, \tag{1.3}$$

where λ is Euler's constant. Thus condition (1.2) is stronger than condition (1.1).

Bourgain already mentionned in his famous paper [1], concerning condition (1.2), that "*... A detailed analysis of previous construction shows that Koksma double logarithmic condition is essentially best possible.*" See Remark p.89. The construction given by Bourgain is one of the several striking examples of application of the entropy criteria he demonstrated in the same paper.

Let $s(n)$ be a nonnegative function of a natural argument. We say that s is submultiplicative if

$$s(nm)\le s(n)s(m),\qquad \forall n,m\ge 1. \tag{1.4}$$

As

$$\sigma_{-1}(n)=\prod_{i=1}^{r}\big(1+\frac{1}{p_i}+\dots\frac{1}{p_i^{\alpha_i}}\big),\qquad n=\prod_{i=1}^{r}p_i^{\alpha_i},$$

Key words and phrases. dilated functions, mean convergence, almost convergence, orthogonal systems, GCD matrix, eigenvalues, quadratic forms.

it is obvious that σ_{-1} is submultiplicative.

We also say that s is bounded in mean if

$$\limsup_{J\to\infty} \frac{1}{J}\sum_{j=1}^{J} s(j) < \infty. \tag{1.5}$$

By Wintner's theorem ([6], p.180) we have

$$\lim_{J\to\infty} \frac{1}{J}\sum_{j=1}^{J} \sigma_{-1}(j) = \sum_{d=1}^{\infty}\frac{1}{d^2}. \tag{1.6}$$

Thus the function σ_{-1} is bounded in mean, whereas the double logarithm function is not.

This difference will permit to put Bourgain's remark under a more precise form. Let $w = \{w_n, n \in \mathbb{Z}\}$ be a sequence of positive reals. Let L^2_w be the associated Sobolev space on the circle, namely the subspace of L^2 consisting with functions f such that

$$\|f\|^2_w := \sum_{n\in\mathbb{Z}} w_n a_n^2(f) < \infty.$$

This is an Hilbert space with scalar product defined by $\langle f, g\rangle = \sum_{n\in\mathbb{Z}} w_n a_n(f)a_n(g)$, $f, g \in L^2_w$.

Theorem 1.2. *Let $w(n)$ be a nonnegative function of a natural argument, which is submultiplicative and bounded in mean. Assume that*

$$w(n) = o(\log\log n).$$

Then there exists a function $f \in L^2_w$ such that the sequence of averages $\frac{1}{n}\sum_{k=1}^{n} f(kx)$, $n = 1, 2, \dots$ fail to converge almost everywhere.

2. Proof

The proof is based on an adaptation to the Sobolev space L^2_w of the method elaborated by Bourgain in [1]. Let $f(x) = \sum_{\ell\in\mathbb{Z}} a_\ell e_\ell$, $a_0 = 0$ and consider the dilation operators $T_j f(x) = f(jx)$. These are positive isometries on L^p, $p \geq 1$, such that $T_j 1 = 1$ for all j, and for all $f \in L^2$

$$\frac{1}{J}\sum_{j\le J} T_j f \xrightarrow{L^2} \int f d\lambda, \qquad J \to \infty,$$

To $f \in L^2$ we associate

$$F_{J,f} = \frac{1}{\sqrt{J}}\sum_{1\le j\le J} g_j T_j f, \qquad (J \geq 1)$$

where $g_1, g_2, \dots$ are i.i.d. standard Gaussian random variables.

Proposition 2.1. *Let $S_n \colon L^2 \to L^2$, $n = 1, 2, \dots$ be continuous operators commuting with T_j on L^2, $S_nT_j = T_jS_n$ for all n and j. Assume that the following property is fulfilled:*

$$\lambda\Big\{\sup_{n\ge1} |S_n(f)| < \infty\Big\} = 1, \qquad \forall f \in L^2_w,$$

then there exists a constant C depending on $\{S_n, n \geq 1\}$ only, such that

$$\sup_{\varepsilon>0} \varepsilon\sqrt{\log N_f(\varepsilon)} \le C \limsup_{J\to\infty} \left(\mathbb{E}\,\|F_{J,f}\|_w\right)^{1/2}, \qquad \forall f \in L^2_w,$$

where $N_f(\varepsilon)$ is the entropy number associated with the set $C_f = \{S_n f, n \geq 1\}$, namely the minimal number of L^2 open balls of radius ε, centered in C_f and enough to cover C_f.

Proof. By the Banach principle, there exists a non-increasing function $C : \mathbb{R}_+ \to \mathbb{R}_+$ such that

$$\forall \varepsilon > 0, \forall g \in L^2_w(\mathbb{T}), \qquad \lambda\Big\{ \sup_n |S_n(g)| \ge \|g\|_w C(\varepsilon) \Big\} \le \varepsilon.$$

Let $0 < \varepsilon < 1/4$. Let $f \in L^2(\mathbb{T})$. Taking $g = F_{J,f}$ and using Fubini's theorem, gives

$$\int_{\mathbb{T}} \mathbb{P}\Big\{ \sup_{n\ge 1} |S_n(F_{J,f})| \ge C(\varepsilon)\|F_{J,f}\|_w \Big\}\, d\lambda \le \varepsilon.$$

It follows that

$$\lambda\left\{ x \in \mathbb{T} : \mathbb{P}\Big\{ \omega : \sup_{n\ge 1} |S_n(F_{J,f}(\omega, .))(x)\ | \ge C(\varepsilon)\|F_{J,f}(\omega, .)\|_w \Big\} \ge \sqrt{\varepsilon} \right\} \le \sqrt{\varepsilon},$$

which is better rewritten under the following form

$$\lambda\left\{ x \in \mathbb{T} : \mathbb{P}\Big\{ \omega : \sup_{n\ge 1} |S_n(F_{J,f}(\omega, .))(x)\ | \le C(\varepsilon)\|F_{J,f}(\omega, .)\|_w \Big\} \ge 1 - \sqrt{\varepsilon} \right\} \ge 1 - \sqrt{\varepsilon}.$$

By Tchebycheff's inequality, $\mathbb{P}\Big\{ \|F_{J,f}\|_w^2 > \mathbb{E}\,\|F_{J,f}\|_w^2/\varepsilon \Big\} \le \varepsilon$. We deduce that the set

$$X_{\varepsilon,J,f} = \Big\{ x \in \mathbb{T} : \mathbb{P}\Big\{ \omega : \sup_{n\ge 1} |S_n(F_{J,f}(\omega, .)(x)| \le C(\varepsilon)\big(\mathbb{E}\,\|F_{J,f}\|_w^2/\varepsilon\big)^{1/2} \Big\} \ge 1 - 2\sqrt{\varepsilon} \Big\}$$

has measure greater than $1-\sqrt{\varepsilon}$. The classical estimate of Gaussian semi-norms implies

$$\forall x \in X_{\varepsilon,J,f}, \qquad \mathbf{E} \sup_{n\ge 1} |S_n(F_{J,f}(\omega,))(x)| \le \frac{4}{(1-2\sqrt{\varepsilon})} \frac{C(\varepsilon)}{\sqrt{\varepsilon}} \Big(\mathbb{E}\,\|F_{J,f}\|_w^2 \Big)^{1/2}.$$

Now let I be a finite set of integers such that $\|S_n(f) - S_m(f)\|_2 \ne 0$, for all distinct elements $m, n \in I$. By the commutation property, $S_n(F_{J,f}) = F_{J,S_n f}$; so that

$$\begin{aligned}
\mathbb{E}\,\big|S_n(F_{J,f}) - S_m(F_{J,f}\big|^2 &= \mathbb{E}\,\big|F_{J,S_n f - S_m f}\big|^2 = \frac{1}{J}\sum_{j\le J} (T_j(S_n f - S_m f))^2 \\
&= \frac{1}{J}\sum_{j\le J} T_j(S_n f - S_m f)^2 \ \to\ \|S_n f - S_m f\|_2^2,
\end{aligned}$$

in L^2 as J tends to infinity. We have used the fact that $(T_j f)^2 = T_j f^2$, if $f \in L^2$. By proceeding by extraction, we can find a partial index $\mathcal{J}$ such that the set

$$A(I) = \left\{ \forall J \in \mathcal{J},\ \forall n, m \in I,\ m \ne n, \quad \frac{\big(\mathbb{E}\,|S_n(F_{J,f}) - S_m(F_{J,f})|^2\big)^{1/2}}{\|S_n(f) - S_m(f)\|_2} \ge \sqrt{1-\varepsilon} \right\},$$

has measure greater that $1 - \sqrt{\varepsilon}$.

Let $J \in \mathcal{J}$, then $\lambda(A(I) \cap X_{\varepsilon,J,f}) \ge 1 - 2\sqrt{\varepsilon} > 0$, and for any $x \in A(I) \cap X_{\varepsilon,J,f}$

$$C(\varepsilon)\Big(\mathbb{E}\,\|F_{J,f}\|_w^2 \Big)^{1/2} \ge \mathbf{E} \sup_{n\ge 1} |S_n(F_{J,f})(x)| \ge \mathbf{E} \sup_{n\in I} S_n(F_{J,f})(x) \ge$$

$$\sqrt{1-\varepsilon}\ \mathbb{E} \sup_{n\in I} Z(S_n(f)) \ge (1 - 2\sqrt{\varepsilon})\ \ \mathbb{E} \sup_{n\in I} Z(S_n(f)).$$

Therefore,

$$\mathbb{E} \sup_{n\in I} Z(S_n(f)) \le C\Big(\limsup_{J\to\infty} \mathbb{E}\,\|F_{J,f}\|_w^2 \Big)^{1/2}.$$

Sudakov's minoration implies

$$\sup_{\rho>0} \rho\sqrt{\log N_f(\rho)} \le C\Big(\limsup_{J\to\infty} \mathbb{E}\,\|F_{J,f}\|_w^2 \Big)^{1/2}.$$

□

We pass to the proof of the theorem. Let $P_1, P_2, \ldots$ denote the sequence of prime numbers. Fix some positive integer s and let d be some other integer such that $2^d \le P_s$. There exists an integer T such that if

$$A_T = \{n = P_1^{\alpha_1} \ldots P_s^{\alpha_s} : 2^T \le n < 2^{T+1},\ \alpha_i \ge 0,\ i = 1, \ldots, s\},$$

then $\sharp(A_{T+d}) \le 2\sharp(A_T)$. Put

$$f = f_T = \frac{1}{\sharp(A_T)^{1/2}} \sum_{n \in A_T} e_n.$$

It follows from Bourgain's proof [1] p.88-89, (or [5] p.239-240 for details) that

$$N\Big(\big(S_{4^i}(f), i \le [\frac{d}{2}]\big), \frac{1}{8}\Big) \ge T.$$

So that

$$\sqrt{\log T} \le C\Big(\limsup_{J\to\infty} \mathbb{E}\,\|F_{J,f}\|_w^2\Big)^{1/2}. \tag{2.1}$$

Now as

$$F_{J,f} = \frac{1}{J^{1/2}} \sum_{j \le J} g_j \frac{1}{\#(A_T)^{1/2}} \sum_{n \in A_T} e_{nj} = \frac{1}{(J\#(A_T))^{1/2}} \sum_{\nu \ge 1} e_\nu \Big(\sum_{\substack{1\le j\le J \\ j|\nu \\ \frac{\nu}{j} \in A_T}} g_j \Big)$$

we have

$$\|F_{J,f}\|_w^2 = \frac{1}{J\#(A_T)} \sum_{\nu \ge 1} w_\nu \Big(\sum_{\substack{1\le j\le J \\ j|\nu \\ \frac{\nu}{j} \in A_T}} g_j \Big)^2.$$

These sums are finite sums. Further,

$$\begin{aligned}
\mathbb{E}\,\|F_{J,f}\|_w^2 &= \frac{1}{J\#(A_T)} \sum_{\nu=1}^{\infty} w_\nu \Big(\sum_{\substack{1\le j\le J \\ j|\nu \\ \frac{\nu}{j} \in A_T}} 1 \Big) = \frac{1}{J\#(A_T)} \sum_{j\le J} \sum_{m \in A_T} w_{mj} \\
&\le \Big(\frac{1}{J} \sum_{j \le J} w_j\Big)\Big(\frac{1}{\#(A_T)} \sum_{m\in A_T} w_m\Big) \\
&\le \Big(\frac{1}{J} \sum_{j \le J} w_j\Big) \max_{m\in A_T} w_m
\end{aligned}$$

Therefore,

$$\limsup_{J\to\infty} \mathbb{E}\,\|F_{J,f}\|_w^2 \le \Big(\limsup_{J\to\infty} \frac{1}{J} \sum_{j\le J} w_j\Big) \max_{m\in A_T} w_m \le M \max_{m\in A_T} w_m,$$

where $M < \infty$ and further

$$\max_{m\in A_T} w_m = o\big(\max_{m\in A_T} \log\log m\big) = o(\log T),$$

by assumption. Consequently

$$\limsup_{J\to\infty} \big(\mathbb{E}\,\|F_{J,f_T}\|_w\big)^{1/2} = o(\sqrt{\log T}). \tag{2.2}$$

But this contradicts (2.1). The theorem is thus proved.

References

[1] Bourgain, J. (1988) Almost sure convergence and bounded entropy, *Israel J. Math.* **63**, 79–95.

[2] Gronwall T.H. (1912) *Some asymptotic expressions in the theory of numbers*, Trans. Am. Math. Soc. **8**, 118–122.

[3] Koksma J.F. (1951) *A diophantine property of summable functions*, J. Indian Math. Soc.(N.S.) **15**, 87–96.

[4] Koksma J.F. (1953) *Estimations de fonctions à l'aide d'intégrales de Lebesgue*, Bull. Soc. Math. Belg. **6**, 4–13.

[5] Weber M. (2009) *Dynamical Systems and Processes*, Eur. Math. Soc. Pub. House, IRMA Lectures in Math. and Theor. Physics **14**.

[6] Wintner A. (1944) *The theory of measure in arithmetical semi-groups*, Wawerly Press, Baltimore, Md.

MICHEL WEBER: IRMA, 10 RUE DU GÉNÉRAL ZIMMER, 67084 STRASBOURG CEDEX, FRANCE
E-mail address: `michel.weber@math.unistra.fr ; m.j.g.weber@gmail.com`